

Table 1.9
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 1999-2008

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.0*	-1.8*	-0.2			
Oral Cavity & Pharynx:	-2.8*	-3.3*	-1.4			
Lip	-	-	-			
Tongue	-3.0*	-2.9*	-2.8			
Salivary gland	-0.7	-4.3*	2.3			
Floor of mouth	-8.7*	-10.3*	-			
Gum & other oral cavity	-4.1	-6.7*	-0.7			
Nasopharynx	-1.6	-1.8	-			
Tonsil	0.3	0.2	-			
Oropharynx	-0.1	0.1	-			
Hypopharynx	-2.5	-1.5	-			
Other oral cavity & pharynx	-	-	-			
Digestive System:	-0.8*	-0.6	-1.1*			
Esophagus	-4.7*	-4.1*	-5.8*			
Stomach	-1.6*	-0.7	-2.3*			
Small intestine	-0.7	-1.1	-0.4			
Colon & Rectum:	-1.8*	-1.8*	-1.9*			
Colon	-2.2*	-2.4*	-2.1*			
Rectum	-0.4	0.3	-1.2			
Anus, anal canal & anorectum	1.8	2.7	1.2			
Liver & intrahep. bile duct:	3.6*	3.8*	2.9			
Liver	3.8*	4.4*	2.2			
Intrahepatic bile duct	0.9	-	-			
Gallbladder	2.4	-	1.7			
Other biliary	4.1*	4.1	3.8*			
Pancreas	0.9	0.5	1.4			
Retroperitoneum	-2.2	-	-			
Peritoneum, omentum & mesentery	-	-	-			
Other digestive system	4.2	-	-			
Respiratory System:	-2.0*	-2.9*	-0.7			
Nose, nasal cavity & middle ear	0.7	-	-			
Larynx	-4.0*	-3.9*	-4.3			
Lung & bronchus	-1.8*	-2.8*	-0.6			
Pleura	-	-	-			
Trachea & other respiratory organs	-	-	-			
Bones & joints	1.9	-0.3	4.2			
Soft tissue (including heart)	-0.1	-2.6	1.9			
Skin (excl. basal & squamous):	0.2	-1.7	1.8			
Melanoma of the skin	1.0	-2.0	-			
Other non-epithelial skin	-0.4	-1.6	1.1			
Breast	0.1	-3.7	0.1			
Breast (<i>in situ</i>)	2.4*	-	2.3*			

- ^a The APC is the Annual Percent Change over the time interval.
SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- * The APC is significantly different from zero (p<.05).
- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.9 - continued
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 1999-2008

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	0.0	-	0.0			
Cervix uteri	-4.3*	-	-4.1*			
Corpus uteri	2.1*	-	2.2*			
Uterus, NOS	4.3	-	4.3			
Ovary ^c	-0.3	-	-0.2			
Vagina	-4.2	-	-3.9			
Vulva	0.5	-	0.7			
Other female genital system	3.3	-	2.7			
Male Genital System:	-2.4*	-2.6*	-			
Prostate	-2.5*	-2.7*	-			
Testis	-1.0	-1.1	-			
Penis	-	-	-			
Other male genital system	-	-	-			
Urinary System:	1.3*	1.6*	0.8			
Urinary bladder	-0.3	0.4	-1.4			
Kidney & renal pelvis	2.9*	3.0*	2.7*			
Ureter	-	-	-			
Other urinary system	-	-	-			
Eye & Orbit	-	-	-			
Brain & Nervous System: ^d	-0.7	-1.3	0.2			
Brain	-0.5	-1.3	0.7			
Cranial nerves & other nervous system	-	-	-			
Endocrine System:	6.1*	5.6*	6.5*			
Thyroid	6.1*	5.9*	6.3*			
Other endocrine & thymus	6.4*	4.6	-			
Lymphoma:	0.8	0.2	1.3*			
Hodgkin lymphoma	1.9*	1.1	2.6			
Non-Hodgkin lymphoma	0.6	0.0	1.1			
Myeloma	0.4	0.5	0.5			
Leukemia:	-1.1	-0.5	-1.6			
Lymphocytic:	-0.5	0.4	-1.8			
Acute lymphocytic	3.8	8.1*	-0.8			
Chronic lymphocytic	-1.2	-0.7	-1.7			
Other lymphocytic	-	-	-			
Myeloid & Monocytic:	-1.7	-1.6	-1.4			
Acute myeloid	-1.9	-2.5	-0.8			
Chronic myeloid	-0.7	0.2	-1.0			
Acute monocytic	-	-	-			
Other myeloid & monocytic	-	-	-			
Other leukemia:	-1.0	-	-			
Other acute leukemia	-	-	-			
Aleukemic, subleukemic & NOS	4.6	-	-			
Kaposi Sarcoma ^e	-4.5*	-4.4*	-			
Mesothelioma ^e	-	-	-			
Ill-defined & unspecified	-3.5*	-5.2*	-2.3			

- ^a The APC is the Annual Percent Change over the time interval.
SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- ^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.
- ^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.
- * The APC is significantly different from zero (p<.05).
- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.